Approccio terapeutico non farmacologico al dolore cronico neuropatico

Stimolazione Cerebrale Non Invasiva - NIBS

Dr med Paolo Rossi

Non Invasive Brain Stimulation

La **NIBS** (*Non-Invasive Brain Stimulation*, ovvero *Stimolazione Cerebrale Non Invasiva*) è un insieme di tecniche utilizzate per modulare l'attività neuronale del cervello senza la necessità di interventi chirurgici o impianti.

TMS (Transcranial Magnetic Stimulation)

utilizza campi magnetici generati da una bobina posizionata sul cuoio capelluto per indurre correnti elettriche nei tessuti cerebrali sottostanti, influenzando l'attività neuronale.

tDCS (Transcranial Direct Current Stimulation)

applica una debole corrente elettrica continua attraverso elettrodi posizionati sullo scalpo per modulare l'eccitabilità corticale (aumentandola o riducendola).

TMS - Stimolazione Magnetica Transcranica

- Single-Pulse TMS – Singolo impulso magnetico
- Paired-Pulse TMS – Coppia di impulsi
- Repetitive TMS (rTMS) – “treni” di impulsi applicati a bassa od alta frequenza

TMS - Stimolazione Magnetica Transcranica

La applicazione di stimoli ripetuti di TMS (rTMS) è in grado di detrminare modificazioni della eccitabilità corticale

$\leq 1\text{Hz}$ ↓ eccitabilità corticale
 $\geq 5\text{Hz}$: ↑ eccitabilità corticale

rTMS - Meccanismo d'azione

Central neuropathic pain

Pain relief

rTMS - Protocolli di Stimolazione

- Area motoria MI controlaterale alla sede del dolore (se bilaterale, emisfero dominante)
- Stimolazione eccitatoria - frequenza 10-20Hz
- Sessioni quotidiane per 5-10 giorni consecutivi

- Responder: pain relief >30%

Contents lists available at [ScienceDirect](https://www.sciencedirect.com)

Clinical Neurophysiology

journal homepage: www.elsevier.com/locate/clinph

Evidence-based guidelines on the therapeutic use of repetitive transcranial magnetic stimulation (rTMS): An update (2014–2018)

Contents

1. Introduction	476	5. Multiple sclerosis	492
2. Pain	477	6. Epilepsy	493
2.1. Motor cortex stimulation in neuropathic pain	477	7. Disorders of consciousness	493
2.2. Other cortical targets in neuropathic pain	479	8. Mild cognitive impairment and Alzheimer's disease	494
2.3. Fibromyalgia and other dysfunctional chronic pain syndromes	480	8.1. HF-rTMS of the left DLPFC	494
2.4. Other pain conditions	480	8.2. HF-rTMS of the precuneus	495
3. Movement disorders	481	8.3. Multisite HF-rTMS	495
3.1. Parkinson's disease: Motor symptoms	481	9. Tinnitus	496
3.2. Parkinson's disease: levodopa-induced dyskinesia	482	9.1. LF-rTMS of the auditory cortex	496
3.3. Parkinson's disease: Depression	483	9.2. HF-rTMS of the left DLPFC	499
3.4. Dystonia	483	10. Depression	500
3.5. Essential tremor	483	10.1. General results	501
3.6. Miscellaneous: Tics and Tourette's syndrome; restless legs syndrome	484	10.2. Novel rTMS protocols to treat depression: Deep HF-rTMS over the left DLPFC	503
4. Stroke	484	10.3. Novel rTMS protocols to treat depression: iTBS over the left DLPFC or cTBS over the right DLPFC	504
4.1. Contralesional LF-rTMS or cTBS at postacute (subacute) stage of limb motor stroke	484	10.4. Novel rTMS protocols to treat depression: Accelerated protocols	506
4.2. Ipsilesional HF-rTMS or iTBS at postacute (subacute) stage of limb motor stroke	486	10.5. Novel rTMS protocols to treat depression: Miscellaneous	506
4.3. Contralesional LF-rTMS at the chronic stage of limb motor stroke	487	10.6. Summary	506
4.4. Ipsilesional HF-rTMS or iTBS at the chronic stage of limb motor stroke	487	11. Schizophrenia	507
4.5. Ipsilesional cTBS at the chronic stage of limb motor stroke	488	11.1. Auditory-verbal hallucinations	507
4.6. Cerebellar target	488	11.2. Negative symptoms	508
4.7. Swallowing and dysphagia	488	12. Substance abuse, addiction and craving	509
4.8. Aphasia	489	12.1. Alcohol craving	509
4.9. Neglect	491	12.2. Nicotine craving	509
		12.3. Methamphetamine or drug craving	510
		12.4. Eating disorders	510
		12.5. Gambling disorders	510
		13. Miscellaneous psychiatric conditions	511
		13.1. Anxiety disorders	511
		13.1.1. Generalized anxiety disorder	511
		13.1.2. Other anxiety disorders	511
		13.2. Post-traumatic stress disorder	511
		13.3. Obsessive compulsive disorder	512
		13.3.1. LF-rTMS of orbitofrontal/prefrontal regions	512
		13.3.2. HF-rTMS of prefrontal regions	513
		13.3.3. LF-rTMS of pre-SMA	513
		13.4. Autism spectrum disorders	514
		13.5. Attention deficit hyperactivity disorder	514
		13.6. Mental retardation	514
		13.7. Functional neurological disorders	514

Evidence-based guidelines on the therapeutic use of repetitive transcranial magnetic stimulation (rTMS): An update (2014–2018)

Table 1
HF-rTMS of M1 contralateral to pain region in neuropathic pain.

Articles	Number of patients	Target, coil type	Control condition	Stimulation frequency and intensity	Number of pulses/session and number of sessions	Significant clinical effects of real versus sham condition	Class of the study
Khedr et al. (2015)	30 patients with malignant neuropathic pain (real: 15; sham: 15)	Hand M1 contralateral to pain, F8c (anteroposterior orientation)	Tilted coil	20 Hz, 80% RMT	2000 pulses, 10 sessions	Reduction of pain score at the end of rTMS protocol (49% on VRS and 37% on VAS), up to 2 weeks after the last session (46% on VRS and 36% on VAS); 87–80% responders (>30% pain relief)	II
Ma et al. (2015)	40 patients with postherpetic neuralgia (real: 20; sham: 20)	Homotopic M1 contralateral to pain region, F8c (anteroposterior orientation)	Tilted coil	10 Hz, 80% RMT	1500 pulses, 10 sessions	Reduction of pain score (17% on VAS), up to 3 months after the last session; 50% responders (>50% pain relief)	II
Attal et al., 2016	32 patients with neuropathic lumbar radicular pain (real: 21; sham: 11)	Hand M1 contralateral to pain, F8c (anteroposterior orientation)	Sham coil	10 Hz, 80% RMT	3000 pulses, 3 sessions	Reduction of pain score at the end of rTMS protocol (#60% on VAS), up to 5 days after the last session (#25% on VAS); 43% responders (>30% pain relief)	II
Nurmikko et al. (2016)	27 patients with neuropathic pain of various origins (crossover)	Homotopic M1 contralateral to pain region or an adjacent motor region, F8c (perpendicular to central sulcus)	Occipital stimulation	10 Hz, 90% RMT	2000 pulses, 5 sessions	Reduction of pain score compared to control condition one week after the last session (9–11% on VAS); 30% responders (>30% pain relief)	II

Pharmacotherapy and non-invasive neuromodulation for neuropathic pain: a systematic review and meta-analysis

Nadia Soliman*, Xavier Moisset*, Michael C Ferraro, Daniel Ciampi de Andrade, Ralf Baron, Joletta Belton†, David L H Bennett, Margarita Calvo, Patrick Dougherty, Ian Gilron, Aki J Hietaharju, Koichi Hosomi, Peter R Kamerman, Harriet Kemp, Elena K Enax-Krumova, Ewan McNicol, Theodore J Price, Srinivasa N Raja, Andrew S C Rice, Blair H Smith, Fiona Talkington†, Andrea Truini, Jan Vollert, Nadine Attal*, Nanna B Finnerup*, Simon Haroutounian*, NeuPSIG Review Update Study Group

	Pain responders (n responders/N total)					Treatment withdrawals (n withdrawals/N total)			
	Active	Placebo	Total patients	Number needed to treat (95% CI)	Susceptibility to bias*	Active	Placebo	Total patients	Number needed to harm (95% CI)
Recommended first-line									
α2δ-ligands	3069/9569	1423/6617	16 186	8.9 (7.4–11.1)	1994	906/9319	332/6866	16185	26.2 (20.4–36.5)
Serotonin noradrenaline reuptake inhibitors	858/2207	364/1493	3700	7.4 (5.6–10.9)	1287	282/2363	61/1567	3930	13.9 (10.9–19.0)
Tricyclic antidepressants	272/723	114/720	1443	4.6 (3.2–7.7)	1728	681/1352	23/671	2023	17.1 (11.4–33.6)
Recommended second-line									
Capsaicin 8% patches	397/1242	214/868	2110	13.16 (7.6–50.8)	NA	11/1288	5/893	2181	1129.3 (135.7–∞)
Capsaicin cream	111/249	68/223	472	6.1 (3.1–∞)	297†	53/495	15/479	974	18.6 (10.6–77.1)
Lidocaine 5% plasters	62/249	33/238	487	14.5 (7.8–108.2)	NA	13/257	10/246	503	178.0 (23.9–∞)
Recommended third-line									
Botulinum toxin type A	102/202	18/171	373	2.7 (1.8–5.1)	1029	2/160	3/157	317	216.3 (23.5–∞)
Repetitive transcranial magnetic stimulation targeting–primary motor cortex	67/207	18/168	375	4.2 (2.3–28.3)	514	2/337	3/299	636	651.6 (34.7–∞)
Opioids	229/613	117/558	1171	5.9 (4.1–10.7)	838	84/781	22/745	1526	15.4 (10.8–24.0)

Data are n/N, unless stated otherwise. NA=not applicable. NNT=number needed to treat. *Refers to the number of patients in a trial who do not respond to treatment that would lead to an NNT>10, considered as the cutoff for reasonable clinical benefit. This calculation is not possible for treatments with NNT>10. The higher the numerical value, the lower the susceptibility to bias. If the susceptibility to bias is less than 400, a new study with fewer than 400 participants with no effect could change the NNT to a level that is not clinically meaningful; however a study with a susceptibility to bias score higher than 400 will not.¹⁴ †Susceptible to reporting bias.

Table 1: Pain response, withdrawals, and susceptibility to reporting bias based on number needed to treat

tDCS - Protocolli di stimolazione

- **Elettrodi:** due elettrodi (anodo e catodo) collegati a un generatore di corrente continua.
- **Montaggio:** l'anodo (positivo) aumenta l'eccitabilità neurale, mentre il catodo (negativo) la riduce.
- **Sede Stimolazione:** M1 o Corteccia Pre-Frontale Dorso-Laterale (DLPFC)
- **Corrente:** tipicamente tra **1 e 2 mA**.
- **Durata:** usualmente **20 minuti** per sessione.
- **Frequenza:** di solito **una seduta al giorno per 5–10 giorni consecutivi**.

a-tDCS of M1

a-tDCS_{UHDS} of M1-S1

a-tDCS_{UHDS} of M1-DLPFC

a-tDCS_{UHDS} of M1-V1*

tDCS - Meccanismo d'azione

Central neuropathic pain

Pain relief

REVIEW

Evidence-Based Guidelines and Secondary Meta-Analysis for the Use of Transcranial Direct Current Stimulation in Neurological and Psychiatric Disorders

Author, Year	Study type	CNP type	Sample (size, sex, age)	tDCS site	Intensity/Current flow time	Duration	Analgesic effect
Bae et al., 2014	Randomized control	PSP	14, 7M, 7F, 45–55	M1	2 mA/20 min	3 weeks	Pain relief by VAS
Hassan et al., 2021	Case report	PSP	1, Gender not, 45	DLPFC	2 mA/20 min	2 weeks	Immediate but transient pain relief by VAS
Molero-Chamizo et al., 2021	3 cases report	PSP	3, 2W:43, 72, 1M:54	M1	1.5 mA/20 min	5 sessions	Significant pain relief by AVAS
Fregni et al., 2006	Randomized control	SCI	17, 14M, 3F, 35.7 ± 13.3	M1	2 mA/20 min	16 days	Significant pain relief by VAS, CGI and PGA
Soler et al., 2010	Randomized control	SCI	39, 30M, 9F, 44.1 ± 11.6	M1	2 mA/20 min	10 sessions	Effective relief for continuous and paroxysmal pain by NRS and BPI
Wrigley et al., 2013	Cross-over	SCI	10, 8M, 2F, 56.1 ± 14.9	M1	2 mA/20 min	5 days	Effective relief for recent pain by NRS
Yoon et al., 2014	Prospective control	SCI	16, 12M, 4F, 44.1 ± 8.6	M1	2 mA/20 min	10 days	Significant pain relief by NRS
Ngernyam et al., 2015	Cross-over	SCI	20, 15M, 5F, 44.5 ± 9.16	M1	2 mA/20 min	1 session	Significant pain relief by NRS
Thibaut et al., 2017	Two-phase randomized control	SCI	Phase I: 33, 24M, 9F, 51.2 ± 12.5 Phase II: 9, 7M, 2F, 49.0 ± 14.38	M1	2 mA/20 min	5 days (Phase I) 10 days (Phase II)	Long-lasting pain relief by VAS
Mori et al., 2010	Randomized, sham-control	MS	19, 8M, 11F, 23–68	motor cortex	2 mA/20 min	5 days	Significant pain relief by VAS and MPQ
Ayache et al., 2016a	Cross-over	MS	16, 3M, 13F, 48.9 ± 10.0	DLPFC	2 mA/20 min	3 days	Significant pain relief by VAS and BPI
González-Zamorano et al., 2021	Randomized control	PD	32, Gender not, Age not	M1	2 mA/20 min	2 weeks (10 sessions)	Pain relief by BPI, KPDPs, PPT, TS and CPM

CNP, central neuropathic pain; PSP, post-stroke pain; CPSP, central post-stroke pain; SCI, spinal cord injury; MS, multiple sclerosis; PD, Parkinson's disease; M, male; F, female; tDCS, transcranial direct current stimulation; RMT, resting motor threshold; mA, milliamperes; min, minute; M1, primary motor cortex; DLPFC, dorsolateral prefrontal cortex; VAS, visual analog scale; AVAS, adaptive visual analog scale; NRS, numerical rating scale; BPI, brief pain inventory; MPQ, McGill pain questionnaire; CGI, clinical global impression; PGA, patient global assessment; PPT, pain pressure threshold; KPDPs, King's Parkinson's disease pain scale; TS, temporal summation; CPM, conditioned pain modulation.

Check for updates

OPEN ACCESS

EDITED BY
 Ma Ke,
 Shanghai Jiao Tong University, China

REVIEWED BY
 Hailong Hou,
 QLD Biotherapeutics Inc.,
 United States
 Angela Maria Sanna,
 ATS Sardegna, Italy

*CORRESPONDENCE
 Xue-Qiang Wang
 wangxueqiang@sus.edu.cn
 Yu-Ling Wang
 wangyul@mail.sysu.edu.cn

†These authors have contributed
 equally to this work and share first

Is transcranial direct current stimulation beneficial for treating pain, depression, and anxiety symptoms in patients with chronic pain? A systematic review and meta-analysis

Yu-Rong Wen^{1,3†}, Jian Shi^{2,3†}, Zheng-Yu Hu^{2,3}, Yang-Yang Lin³,
 You-Tian Lin^{3,4}, Xue Jiang¹, Rui Wang¹, Xue-Qiang Wang^{1,5*}
 and Yu-Ling Wang^{3*}

A

B

FIGURE 2

Forest plot showing overall effect sizes (Hedges' g) of real tDCS on pain intensity within studies. These plots show the pooled SMD (large diamond shape) and I^2 resulting from the meta-analysis. **(A)** Pain scores from short-term data. **(B)** Pain scores from medium-term data. CI, confidence interval; SMD, standard mean difference; tDCS, transcranial direct current stimulation; a, motor cortex; b, dorsolateral prefrontal cortex; c, operculo-insular cortex.

Warnings

rTMS

- Epilessia non controllata
- Pacemaker, neurostimolatori
- Clip Metalliche
- Epilessia

tDCS

- Epilessia non controllata
- lesioni cutanee nella sede degli elettrodi
- pacemaker, neurostimolatori

rTMS vs tDCS

	rTMS	tDCS
Facilità uso	<u>+</u>	+
Tollerabilità	+	++
Costi	\$\$	\$
Rimborsabilità	✗	✗

Conclusioni

- rTMS e TDCS sono in grado di rimodulare l'eccitabilità corticale mediando plasticità corticale, neurotrasmettitori SNC e regolando il rilascio di citochine pro-infiammatorie
- Efficacia sul dolore neuropatico più definita per rTMS
- Profilo di sicurezza alto
 - CAVE: epilessia, device impiantabili, clips metalliche (psicosi attive?)
- Costi Device

Grazie per l'ascolto!

www.rehaticino.ch